

Multil-level Weiner-Hopf Monte-Carlo simulation for Lévy processes

A. Ferreiro-Castilla, A.E. Kyprianou, R. Scheichl, G. Suryanarayana

Department of Mathematical Sciences, University of Bath

Motivation

- **Lévy process.** A (one dimensional) process with stationary and independent increments which has paths which are right continuous with left limits and therefore includes Brownian motion with drift, compound Poisson processes, stable processes amongst many others).
- A popular (**and often criticised**) model in mathematical finance for the evolution of a risky asset is

$$S_t := e^{X_t}, t \geq 0$$

where $\{X_t : t \geq 0\}$ is a Lévy process.

- Barrier options: The value of up-and-out barrier option with expiry date T and barrier b is typically priced as

$$\mathbb{E}_s(f(X_1)\mathbf{1}_{\{\bar{X}_1 \leq b\}})$$

where $\bar{X}_1 = \sup_{u \leq 1} X_u$, f is some nice function.

- One is fundamentally interested in the joint distribution


$$\mathbb{P}(X_1 \in dx, \bar{X}_1 \in dy)$$

for any Lévy process (X, \mathbb{P}) .

Original WHMC method: Kuznetsov et al. (2011)


Original WHMC method: Kuznetsov et al. (2011)

- Consider a Poisson process with arrival rate n . Denote by τ_1, τ_2, \dots the arrival times.



Original WHMC method: Kuznetsov et al. (2011)

- Consider a Poisson process with arrival rate n . Denote by τ_1, τ_2, \dots the arrival times.



- Note that τ_n is the sum of n i.i.d exponential random variables, each with mean $1/n$. We could therefore write


$$\tau_n = \sum_{i=1}^n \frac{1}{n} e^{(i)},$$

where $e^{(i)}$ are i.i.d. exponential random variables with unit mean. Hence by the SLLN

$$\tau_n \rightarrow 1 \quad \text{almost surely.}$$

Original WHMC method: Kuznetsov et al. (2011)

- Consider a Poisson process with arrival rate n . Denote by τ_1, τ_2, \dots the arrival times.



- Note that τ_n is the sum of n i.i.d exponential random variables, each with mean $1/n$. We could therefore write

$$\tau_n = \sum_{i=1}^n \frac{1}{n} e^{(i)},$$

where $e^{(i)}$ are i.i.d. exponential random variables with unit mean. Hence by the SLLN

$$\tau_n \rightarrow 1 \quad \text{almost surely.}$$

- Hence for a suitably large n , we have in distribution

$$(X_{\tau_n}, \bar{X}_{\tau_n}) \simeq (X_1, \bar{X}_1).$$

Indeed since 1 is not a jump time with probability 1, we have that $(X_{\tau_n}, \bar{X}_{\tau_n}) \rightarrow (X_1, \bar{X}_1)$ almost surely as $n \rightarrow \infty$.

Original WHMC method: Kuznetsov et al. (2011)

Original WHMC method: Kuznetsov et al. (2011)

- A reformulation of the Wiener-Hopf factorization states that

$$X_{\mathbf{e}_q} \stackrel{d}{=} S_q + I_q$$


where S_q is independent of I_q and they are respectively equal in distribution to $\overline{X}_{\mathbf{e}_q}$ and $\underline{X}_{\mathbf{e}_q}$. Here $\underline{X}_t = \inf_{s \leq t} X_s$.

Original WHMC method: Kuznetsov et al. (2011)

- A reformulation of the Wiener-Hopf factorization states that

$$X_{e_q} \stackrel{d}{=} S_q + I_q$$

where S_q is independent of I_q and they are respectively equal in distribution to \bar{X}_{e_q} and \underline{X}_{e_q} . Here $\underline{X}_t = \inf_{s \leq t} X_s$.



Original WHMC method: Kuznetsov et al. (2011)


- A reformulation of the Wiener-Hopf factorization states that

$$X_{\mathbf{e}_q} \stackrel{d}{=} S_q + I_q$$

where S_q is independent of I_q and they are respectively equal in distribution to $\overline{X}_{\mathbf{e}_q}$ and $\underline{X}_{\mathbf{e}_q}$.

- Taking advantage of the above, the fact that X has stationary and independent increments and the fact that, as a time, τ_n can be seen as the sum of independent exponential time periods we have the following:

Original WHMC method: Kuznetsov et al. (2011)



Original WHMC method: Kuznetsov et al. (2011)

- For all $n \in \{1, 2, \dots\}$ and $n > 0$,

$$(X_{\tau_n}, \bar{X}_{\tau_n}) \stackrel{d}{=} (V_n, J_n)$$

where

$$V_n = \sum_{j=1}^n \{S_n^{(j)} + I_n^{(j)}\}$$

$$J_n := \bigvee_{i=0}^{n-1} (V_i + S_n^{(i+1)}).$$

Here, $V_0 = S_n^{(0)} = I_n^{(0)} = 0$, $\{S_n^{(j)} : j \geq 1\}$ are an i.i.d. sequence of random variables with common distribution equal to that of \bar{X}_{e_n} and $\{I_n^{(j)} : j \geq 1\}$ are another i.i.d. sequence of random variable with common distribution equal to that of \underline{X}_{e_n} .

Original WHMC method: Kuznetsov et al. (2011)

- For all $n \in \{1, 2, \dots\}$ and $n > 0$,

$$(X_{\tau_n}, \bar{X}_{\tau_n}) \stackrel{d}{=} (V_n, J_n)$$

where

$$V_n = \sum_{j=1}^n \{S_n^{(j)} + I_n^{(j)}\}$$

$$J_n := \bigvee_{i=0}^{n-1} (V_i + S_n^{(i+1)}).$$

Here, $V_0 = S_n^{(0)} = I_n^{(0)} = 0$, $\{S_n^{(j)} : j \geq 1\}$ are an i.i.d. sequence of random variables with common distribution equal to that of \bar{X}_{e_n} and $\{I_n^{(j)} : j \geq 1\}$ are another i.i.d. sequence of random variable with common distribution equal to that of \underline{X}_{e_n} .

- $(V_n, J_n) \xrightarrow{n \uparrow \infty} (X_1, \bar{X}_1)$ in distribution.

Original WHMC method: Kuznetsov et al. (2011)

Original WHMC method: Kuznetsov et al. (2011)

- Sample repeatedly and independently from the distribution \overline{X}_{e_n} and \underline{X}_{e_n} and then construct m independent versions of the variables V_n and J_n , say

$$\{V_n^{(i)} : i = 1, \dots, m\} \text{ and } \{J_n^{(i)} : i = 1, \dots, m\}.$$

Original WHMC method: Kuznetsov et al. (2011)

- Sample repeatedly and independently from the distribution \overline{X}_{e_n} and \underline{X}_{e_n} and then construct m independent versions of the variables V_n and J_n , say

$$\{V_n^{(i)} : i = 1, \dots, m\} \text{ and } \{J_n^{(i)} : i = 1, \dots, m\}.$$

- Then

$$\mathbb{E}(F(X_1, \overline{X}_1)) \simeq \mathbb{E}(F(X_{\tau_n}, \overline{X}_{\tau_n})) = \mathbb{E}(F(V_n, J_n)) \simeq \frac{1}{m} \sum_{i=1}^m F(V_n^{(i)}, J_n^{(i)}) =: \widehat{F}_{MC}^{n,m}.$$

Original WHMC method: Kuznetsov et al. (2011)

- Sample repeatedly and independently from the distribution \overline{X}_{e_n} and \underline{X}_{e_n} and then construct m independent versions of the variables V_n and J_n , say

$$\{V_n^{(i)} : i = 1, \dots, m\} \text{ and } \{J_n^{(i)} : i = 1, \dots, m\}.$$

- Then

$$\mathbb{E}(F(X_1, \overline{X}_1)) \simeq \mathbb{E}(F(X_{\tau_n}, \overline{X}_{\tau_n})) = \mathbb{E}(F(V_n, J_n)) \simeq \frac{1}{m} \sum_{i=1}^m F(V_n^{(i)}, J_n^{(i)}) =: \widehat{F}_{MC}^{n,m}.$$

- Sampling from \overline{X}_{e_n} and \underline{X}_{e_n} is generally impossible for a given Lévy process, but not for a 10 parameter family of processes known as Kuznetsov's β -class (ask me afterwards if interested in the details!).

Numerical Analysis

Henceforth assume that

Numerical Analysis

Henceforth assume that

- $F : \mathbb{R} \times [0, \infty) \mapsto [0, \infty)$ is Lipschitz continuous with coefficient 1.

Numerical Analysis

Henceforth assume that

- $F : \mathbb{R} \times [0, \infty) \mapsto [0, \infty)$ is Lipschitz continuous with coefficient 1.
- Our underlying Lévy process satisfies

$$\int_{|x| \geq 1} x^2 \Pi(dx) < \infty,$$

where Π is its associated jump measure (finite second moments).

Numerical Analysis

Henceforth assume that

- $F : \mathbb{R} \times [0, \infty) \mapsto [0, \infty)$ is Lipschitz continuous with coefficient 1.
- Our underlying Lévy process satisfies

$$\int_{|x| \geq 1} x^2 \Pi(dx) < \infty,$$

where Π is its associated jump measure (finite second moments).

■ Notation.

- Write $a \lesssim b$ for two positive quantities a and b , if a/b is uniformly bounded (independent of n , M , or any other parameters).
- Write $F^{n,(i)} := F(V_n^{(i)}, J_n^{(i)})$ for the i -th sample of $F^n := F(V_n, J_n)$ (using the Wiener-Hopf random walk).
- Define the **mean square error** as

$$e(\widehat{F}_{MC}^{n,m})^2 := \mathbb{E}[(\widehat{F}_{MC}^{n,m} - \mathbb{E}[F(X_1, \overline{X}_1)])^2] = m^{-1} \mathbb{V}(F^n) + (\mathbb{E}[F^n - F(X_1, \overline{X}_1)])^2$$

Numerical Analysis

Henceforth assume that

- $F : \mathbb{R} \times [0, \infty) \mapsto [0, \infty)$ is Lipschitz continuous with coefficient 1.
- Our underlying Lévy process satisfies

$$\int_{|x| \geq 1} x^2 \Pi(dx) < \infty,$$

where Π is its associated jump measure (finite second moments).

■ Notation.

- Write $a \lesssim b$ for two positive quantities a and b , if a/b is uniformly bounded (independent of n , M , or any other parameters).
- Write $F^{n,(i)} := F(V_n^{(i)}, J_n^{(i)})$ for the i -th sample of $F^n := F(V_n, J_n)$ (using the Wiener-Hopf random walk).
- Define the **mean square error** as

$$e(\widehat{F}_{MC}^{n,m})^2 := \mathbb{E}[(\widehat{F}_{MC}^{n,m} - \mathbb{E}[F(X_1, \overline{X}_1)])^2] = m^{-1} \mathbb{V}(F^n) + (\mathbb{E}[F^n - F(X_1, \overline{X}_1)])^2$$

- Then we have the following convergence/complexity theorem ...

Numerical Analysis WHMC

Theorem (Single-level WHMC)

Assume that $\exists \alpha > 0$ s.t.

- (i) $\mathbb{E}[|F^n - F(X_t, \bar{X}_t)|] \lesssim n^{-\alpha}$ and
- (ii) $\mathbb{E}[\mathcal{C}(F^n)] \lesssim n$ (where $\mathcal{C}(F^n)$ is the cost to compute a single sample from F^n)

Then, $\forall \nu \in \mathbb{N} \exists n, M \in \mathbb{N}$ s.t.

$$\mathbb{E} \left[\mathcal{C}(\hat{F}_{MC}^{n,M}) \right] \lesssim \nu \quad \text{and } L^2 \text{ error } e(\hat{F}_{MC}^{n,m}) \lesssim \nu^{-\frac{1}{2+1/\alpha}}.$$

Numerical Analysis WHMC

Theorem (Single-level WHMC)

Assume that $\exists \alpha > 0$ s.t.

- (i) $\mathbb{E}[|F^n - F(X_t, \bar{X}_t)|] \lesssim n^{-\alpha}$ and
- (ii) $\mathbb{E}[\mathcal{C}(F^n)] \lesssim n$ (where $\mathcal{C}(F^n)$ is the cost to compute a single sample from F^n)

Then, $\forall \nu \in \mathbb{N} \exists n, M \in \mathbb{N}$ s.t.

$$\mathbb{E} \left[\mathcal{C}(\hat{F}_{MC}^{n,M}) \right] \lesssim \nu \quad \text{and } L^2 \text{ error } e(\hat{F}_{MC}^{n,m}) \lesssim \nu^{-\frac{1}{2+1/\alpha}}.$$

- For Kuznetsov's β -class of Lévy processes also Assumption (ii) holds.

Numerical Analysis WHMC

Theorem (Single-level WHMC)

Assume that $\exists \alpha > 0$ s.t.

- (i) $\mathbb{E}[|F^n - F(X_t, \bar{X}_t)|] \lesssim n^{-\alpha}$ and
- (ii) $\mathbb{E}[\mathcal{C}(F^n)] \lesssim n$ (where $\mathcal{C}(F^n)$ is the cost to compute a single sample from F^n)

Then, $\forall \nu \in \mathbb{N} \exists n, M \in \mathbb{N}$ s.t.

$$\mathbb{E} \left[\mathcal{C}(\hat{F}_{MC}^{n,M}) \right] \lesssim \nu \quad \text{and } L^2 \text{ error } e(\hat{F}_{MC}^{n,m}) \lesssim \nu^{-\frac{1}{2+1/\alpha}}.$$

- For Kuznetsov's β -class of Lévy processes also Assumption (ii) holds.

Using the forthcoming analysis we shall shortly present, it will turn out that:

- when X has paths of unbounded variation, $\alpha = \frac{1}{4} \Rightarrow \mathcal{O}(\nu^{-\frac{1}{6}})$ convergence!
- when X has paths of bounded variation, $\alpha = \frac{1}{2} \Rightarrow \mathcal{O}(\nu^{-\frac{1}{4}})$ convergence!

Numerical Analysis WHMC

Theorem (Single-level WHMC)

Assume that $\exists \alpha > 0$ s.t.

- (i) $\mathbb{E}[|F^n - F(X_t, \bar{X}_t)|] \lesssim n^{-\alpha}$ and
- (ii) $\mathbb{E}[\mathcal{C}(F^n)] \lesssim n$ (where $\mathcal{C}(F^n)$ is the cost to compute a single sample from F^n)

Then, $\forall \nu \in \mathbb{N} \exists n, M \in \mathbb{N}$ s.t.

$$\mathbb{E} \left[\mathcal{C}(\hat{F}_{MC}^{n,M}) \right] \lesssim \nu \quad \text{and } L^2 \text{ error } e(\hat{F}_{MC}^{n,m}) \lesssim \nu^{-\frac{1}{2+1/\alpha}}.$$

- For Kuznetsov's β -class of Lévy processes also Assumption (ii) holds.

Using the forthcoming analysis we shall shortly present, it will turn out that:

- when X has paths of unbounded variation, $\alpha = \frac{1}{4} \Rightarrow \mathcal{O}(\nu^{-\frac{1}{6}})$ convergence!
- when X has paths of bounded variation, $\alpha = \frac{1}{2} \Rightarrow \mathcal{O}(\nu^{-\frac{1}{4}})$ convergence!

The best one can hope for with such Monte-Carlo schemes is an $\mathcal{O}(\nu^{-\frac{1}{2}})$ convergence.

Multi-level Wiener-Hopf Monte-Carlo [Heinrich, 2001], [Giles, 2007], ...

- Computational gains from exploiting the telescopic sum

$$\mathbb{E}[F^{n_L}] = \mathbb{E}[F^{n_0}] + \sum_{\ell=1}^L \mathbb{E}[F^{n_\ell} - F^{n_{\ell-1}}],$$

where $n_\ell = 2^\ell n_0$, $\ell = 1, \dots, L$, for some small $n_0 \in \mathbb{N}$.

Multi-level Wiener-Hopf Monte-Carlo [Heinrich, 2001], [Giles, 2007], ...

- Computational gains from exploiting the telescopic sum

$$\mathbb{E}[F^{n_L}] = \mathbb{E}[F^{n_0}] + \sum_{\ell=1}^L \mathbb{E}[F^{n_\ell} - F^{n_{\ell-1}}],$$

where $n_\ell = 2^\ell n_0$, $\ell = 1, \dots, L$, for some small $n_0 \in \mathbb{N}$.

- Suggesting the multilevel estimator

$$\widehat{F}_{ML}^{n_0, L, \{M_\ell\}} := \frac{1}{M_0} \sum_{i=1}^{M_0} F^{n_0, (i)} + \sum_{\ell=1}^L \frac{1}{M_\ell} \sum_{i=1}^{M_\ell} (F^{n_\ell, (i)} - F^{n_{\ell-1}, (i)}).$$

- Here it is **very important** that $F^{n_{\ell-1}}$ can be obtained from F^{n_ℓ} by a “deterministic” transformation of the random variables used to obtain F^{n_ℓ} .

Multi-level Wiener-Hopf Monte-Carlo [Heinrich, 2001], [Giles, 2007], ...

- Computational gains from exploiting the telescopic sum

$$\mathbb{E}[F^{n_L}] = \mathbb{E}[F^{n_0}] + \sum_{\ell=1}^L \mathbb{E}[F^{n_\ell} - F^{n_{\ell-1}}],$$

where $n_\ell = 2^\ell n_0$, $\ell = 1, \dots, L$, for some small $n_0 \in \mathbb{N}$.

- Suggesting the multilevel estimator

$$\widehat{F}_{ML}^{n_0, L, \{M_\ell\}} := \frac{1}{M_0} \sum_{i=1}^{M_0} F^{n_0, (i)} + \sum_{\ell=1}^L \frac{1}{M_\ell} \sum_{i=1}^{M_\ell} (F^{n_\ell, (i)} - F^{n_{\ell-1}, (i)}).$$

- Here it is **very important** that $F^{n_{\ell-1}}$ can be obtained from F^{n_ℓ} by a “deterministic” transformation of the random variables used to obtain F^{n_ℓ} .
- A little algebra again reveals that the means square error satisfies

$$e(\widehat{F}_{ML}^{n_0, L, \{M_\ell\}})^2 = \frac{1}{M_0} \mathbb{V}(F^{n_0}) + \sum_{\ell=1}^L \frac{1}{M_\ell} \mathbb{V}(F^{n_\ell} - F^{n_{\ell-1}}) + (\mathbb{E}[F^n - F(X_1, \bar{X}_1)])^2.$$

Multi-level Wiener-Hopf Monte-Carlo [Heinrich, 2001], [Giles, 2007], ...

- Computational gains from exploiting the telescopic sum

$$\mathbb{E}[F^{n_L}] = \mathbb{E}[F^{n_0}] + \sum_{\ell=1}^L \mathbb{E}[F^{n_\ell} - F^{n_{\ell-1}}],$$

where $n_\ell = 2^\ell n_0$, $\ell = 1, \dots, L$, for some small $n_0 \in \mathbb{N}$.

- Suggesting the multilevel estimator

$$\widehat{F}_{ML}^{n_0, L, \{M_\ell\}} := \frac{1}{M_0} \sum_{i=1}^{M_0} F^{n_0, (i)} + \sum_{\ell=1}^L \frac{1}{M_\ell} \sum_{i=1}^{M_\ell} (F^{n_\ell, (i)} - F^{n_{\ell-1}, (i)}).$$

- Here it is **very important** that $F^{n_{\ell-1}}$ can be obtained from F^{n_ℓ} by a “deterministic” transformation of the random variables used to obtain F^{n_ℓ} .
- A little algebra again reveals that the means square error satisfies

$$e(\widehat{F}_{ML}^{n_0, L, \{M_\ell\}})^2 = \frac{1}{M_0} \mathbb{V}(F^{n_0}) + \sum_{\ell=1}^L \frac{1}{M_\ell} \mathbb{V}(F^{n_\ell} - F^{n_{\ell-1}}) + (\mathbb{E}[F^{n_L} - F(X_1, \overline{X}_1)])^2.$$

- See also [Dereich, Heidenreich, 2011], [Dereich, 2011], [Giles, Xia, 2012].

Poisson thinning and multilevel WHMC

Poisson thinning and multilevel WHMC



- In the WHMC method how do we introduce "levels"?

Poisson thinning and multilevel WHMC

- In the WHMC method how do we introduce "levels"?
- Recall also that it is **crucial** to have a Poisson process for the time randomisations on **all** levels! How do we sample on two consecutive levels?

Poisson thinning and multilevel WHMC

- In the WHMC method how do we introduce "levels"?
- Recall also that it is **crucial** to have a Poisson process for the time randomisations on **all** levels! How do we sample on two consecutive levels?
- Suppose the "level ℓ " grid is based on a Poisson process of rate n_ℓ . Then by **tossing a coin** and rejecting arrivals with probability $1/2$ we end up with a Poisson process of rate $n_{\ell-1}$: our new coarser "level $\ell - 1$ " Poisson grid. (Not a new idea! Also used by [Glasserman, Merener, 2003], [Giles, Xia, 2012], ...)



Numerical Analysis (multilevel case)

Theorem (Multilevel WHMC)

Assume $\exists \alpha, \beta > 0$ with $\alpha \geq \frac{1}{2} \max\{\beta, 1\}$ such that

- (i) $|\mathbb{E}[F^{n_\ell} - F(X_t, \bar{X}_t)]| \lesssim n_\ell^{-\alpha}$
- (ii) $\mathbb{V}[F^{n_\ell} - F^{n_{\ell-1}}] \lesssim n_\ell^{-\beta}$
- (iii) $\mathbb{E}[C_{n_\ell}] \lesssim n_\ell$.

Then, $\forall \nu \in \mathbb{N} \exists L$ and $\{M_\ell\}_{\ell=0}^L$ s.t. $\mathbb{E}\left[\mathcal{C}\left(\widehat{F}_{\text{ML}}^{n_0, L, \{M_\ell\}}\right)\right] \lesssim \nu$ and L^2 error

$$e\left(\widehat{F}_{\text{ML}}^{n_0, L, \{M_\ell\}}\right) \lesssim \begin{cases} \nu^{-\frac{1}{2}} & \text{if } \beta > 1, \\ \nu^{-\frac{1}{2}} \log^2 \nu & \text{if } \beta = 1, \\ \nu^{-\frac{1}{2+(1-\beta)/\alpha}} & \text{if } \beta < 1. \end{cases}$$

Numerical Analysis (multilevel case)

Theorem (Multilevel WHMC)

Assume $\exists \alpha, \beta > 0$ with $\alpha \geq \frac{1}{2} \max\{\beta, 1\}$ such that

- (i) $|\mathbb{E}[F^{n_\ell} - F(X_t, \bar{X}_t)]| \lesssim n_\ell^{-\alpha}$
- (ii) $\mathbb{V}[F^{n_\ell} - F^{n_{\ell-1}}] \lesssim n_\ell^{-\beta}$
- (iii) $\mathbb{E}[C_{n_\ell}] \lesssim n_\ell$.

Then, $\forall \nu \in \mathbb{N} \exists L$ and $\{M_\ell\}_{\ell=0}^L$ s.t. $\mathbb{E}\left[\mathcal{C}\left(\widehat{F}_{\text{ML}}^{n_0, L, \{M_\ell\}}\right)\right] \lesssim \nu$ and L^2 error

$$e\left(\widehat{F}_{\text{ML}}^{n_0, L, \{M_\ell\}}\right) \lesssim \begin{cases} \nu^{-\frac{1}{2}} & \text{if } \beta > 1, \\ \nu^{-\frac{1}{2}} \log^2 \nu & \text{if } \beta = 1, \\ \nu^{-\frac{1}{2 + (1-\beta)/\alpha}} & \text{if } \beta < 1. \end{cases}$$

- For Kuznetsov's β -class of Lévy processes also Assumption (iii) holds.
- Recall that $\alpha = (1/4)1/2$ for (un)bounded variation paths & shortly we shall see $\beta = 1/2 \Rightarrow (\mathcal{O}(\nu^{-\frac{1}{4}})) \mathcal{O}(\nu^{-\frac{1}{3}})$.
- Compare with former (single-level) rates $(\mathcal{O}(\nu^{-\frac{1}{6}})) \mathcal{O}(\nu^{-\frac{1}{4}})$.

Numerical analysis MLWH

Remains to verify the assumptions (i) $|\mathbb{E}[F^{n_\ell} - F(X_t, \bar{X}_t)]| \lesssim n_\ell^{-\alpha}$ and (ii) $\mathbb{V}[F^{n_\ell} - F^{n_{\ell-1}}] \lesssim n_\ell^{-\beta}$.

- Recalling that F is Lipschitz

$$\begin{aligned} \mathbb{V}(F^{n_\ell} - F^{n_{\ell-1}}) &= \mathbb{V}(F(X_{\tau_{n_\ell}}, \bar{X}_{\tau_{n_\ell}}) - F(X_{\tau_{n_{\ell-1}}}, \bar{X}_{\tau_{n_{\ell-1}}})) \\ &\leq \mathbb{E}[(X_{\tau_{n_\ell}} - X_{\tau_{n_{\ell-1}}})^2] + \mathbb{E}[(\bar{X}_{\tau_{n_\ell}} - \bar{X}_{\tau_{n_{\ell-1}}})^2] \end{aligned}$$

Numerical analysis MLWH

Remains to verify the assumptions (i) $|\mathbb{E}[F^{n_\ell} - F(X_t, \bar{X}_t)]| \lesssim n_\ell^{-\alpha}$ and (ii) $\mathbb{V}[F^{n_\ell} - F^{n_{\ell-1}}] \lesssim n_\ell^{-\beta}$.

- Recalling that F is Lipschitz

$$\begin{aligned} \mathbb{V}(F^{n_\ell} - F^{n_{\ell-1}}) &= \mathbb{V}(F(X_{\tau_{n_\ell}}, \bar{X}_{\tau_{n_\ell}}) - F(X_{\tau_{n_{\ell-1}}}, \bar{X}_{\tau_{n_{\ell-1}}})) \\ &\leq \mathbb{E}[(X_{\tau_{n_\ell}} - X_{\tau_{n_{\ell-1}}})^2] + \mathbb{E}[(\bar{X}_{\tau_{n_\ell}} - \bar{X}_{\tau_{n_{\ell-1}}})^2] \end{aligned}$$

- **[Unbounded variation case]:** Working with our Lévy process as a Markovian semi-martingale we get for s and t random times independent of X ,

$$(1) \quad \mathbb{E}[(X_t - X_s)^2] = \mathbb{V}(X_1)\mathbb{E}[|t - s|] + \mathbb{E}[X_1]^2\mathbb{E}[(t - s)^2]$$

$$(2) \quad \mathbb{E}[(\bar{X}_t - \bar{X}_s)^2] \leq 16\mathbb{V}(X_1)\mathbb{E}[|t - s|] + 2(\max\{\mathbb{E}[X_1], 0\})^2\mathbb{E}[(t - s)^2],$$

[Better estimates for bounded variation case!]

Numerical analysis MLWH

Remains to verify the assumptions (i) $|\mathbb{E}[F^{n_\ell} - F(X_t, \bar{X}_t)]| \lesssim n_\ell^{-\alpha}$ and (ii) $\mathbb{V}[F^{n_\ell} - F^{n_\ell-1}] \lesssim n_\ell^{-\beta}$.

- Recalling that F is Lipschitz

$$\begin{aligned} \mathbb{V}(F^{n_\ell} - F^{n_\ell-1}) &= \mathbb{V}(F(X_{\tau_{n_\ell}}, \bar{X}_{\tau_{n_\ell}}) - F(X_{\tau_{n_\ell-1}}, \bar{X}_{\tau_{n_\ell-1}})) \\ &\leq \mathbb{E}[(X_{\tau_{n_\ell}} - X_{\tau_{n_\ell-1}})^2] + \mathbb{E}[(\bar{X}_{\tau_{n_\ell}} - \bar{X}_{\tau_{n_\ell-1}})^2] \end{aligned}$$

- **[Unbounded variation case]:** Working with our Lévy process as a Markovian semi-martingale we get for s and t random times independent of X ,

$$(1) \quad \mathbb{E}[(X_t - X_s)^2] = \mathbb{V}(X_1)\mathbb{E}[|t - s|] + \mathbb{E}[X_1]^2\mathbb{E}[(t - s)^2]$$

$$(2) \quad \mathbb{E}[(\bar{X}_t - \bar{X}_s)^2] \leq 16\mathbb{V}(X_1)\mathbb{E}[|t - s|] + 2(\max\{\mathbb{E}[X_1], 0\})^2\mathbb{E}[(t - s)^2],$$

[Better estimates for bounded variation case!]

- Finally, recalling that τ_{n_ℓ} is a (n_ℓ, n_ℓ) -gamma distribution with mean 1,

$$\mathbb{E}[(\tau_{n_\ell} - 1)^2] \lesssim n_\ell^{-1} \quad \text{and} \quad \mathbb{E}[|\tau_{n_\ell} - 1|] \lesssim n_\ell^{-\frac{1}{2}}.$$

Numerical analysis MLWH

Remains to verify the assumptions (i) $|\mathbb{E}[F^{n_\ell} - F(X_t, \bar{X}_t)]| \lesssim n_\ell^{-\alpha}$ and (ii) $\mathbb{V}[F^{n_\ell} - F^{n_{\ell-1}}] \lesssim n_\ell^{-\beta}$.

- Recalling that F is Lipschitz

$$\begin{aligned} \mathbb{V}(F^{n_\ell} - F^{n_{\ell-1}}) &= \mathbb{V}(F(X_{\tau_{n_\ell}}, \bar{X}_{\tau_{n_\ell}}) - F(X_{\tau_{n_{\ell-1}}}, \bar{X}_{\tau_{n_{\ell-1}}})) \\ &\leq \mathbb{E}[(X_{\tau_{n_\ell}} - X_{\tau_{n_{\ell-1}}})^2] + \mathbb{E}[(\bar{X}_{\tau_{n_\ell}} - \bar{X}_{\tau_{n_{\ell-1}}})^2] \end{aligned}$$

- [Unbounded variation case]:** Working with our Lévy process as a Markovian semi-martingale we get for s and t random times independent of X ,

$$(1) \quad \mathbb{E}[(X_t - X_s)^2] = \mathbb{V}(X_1)\mathbb{E}[|t - s|] + \mathbb{E}[X_1]^2\mathbb{E}[(t - s)^2]$$

$$(2) \quad \mathbb{E}[(\bar{X}_t - \bar{X}_s)^2] \leq 16\mathbb{V}(X_1)\mathbb{E}[|t - s|] + 2(\max\{\mathbb{E}[X_1], 0\})^2\mathbb{E}[(t - s)^2],$$

[Better estimates for bounded variation case!]

- Finally, recalling that τ_{n_ℓ} is a (n_ℓ, n_ℓ) -gamma distribution with mean 1,

$$\mathbb{E}[(\tau_{n_\ell} - 1)^2] \lesssim n_\ell^{-1} \quad \text{and} \quad \mathbb{E}[|\tau_{n_\ell} - 1|] \lesssim n_\ell^{-\frac{1}{2}}.$$

- All together we get $\mathbb{V}(F^{n_\ell} - F^{n_{\ell-1}}) \lesssim n_\ell^{-\frac{1}{2}}$, and so $\beta = \frac{1}{2}$.

Numerical analysis MLWH

Remains to verify the assumptions (i) $|\mathbb{E}[F^{n_\ell} - F(X_t, \bar{X}_t)]| \lesssim n_\ell^{-\alpha}$ and (ii) $\mathbb{V}[F^{n_\ell} - F^{n_{\ell-1}}] \lesssim n_\ell^{-\beta}$.

- Recalling that F is Lipschitz

$$\begin{aligned} \mathbb{V}(F^{n_\ell} - F^{n_{\ell-1}}) &= \mathbb{V}(F(X_{\tau_{n_\ell}}, \bar{X}_{\tau_{n_\ell}}) - F(X_{\tau_{n_{\ell-1}}}, \bar{X}_{\tau_{n_{\ell-1}}})) \\ &\leq \mathbb{E}[(X_{\tau_{n_\ell}} - X_{\tau_{n_{\ell-1}}})^2] + \mathbb{E}[(\bar{X}_{\tau_{n_\ell}} - \bar{X}_{\tau_{n_{\ell-1}}})^2] \end{aligned}$$

- [Unbounded variation case]:** Working with our Lévy process as a Markovian semi-martingale we get for s and t random times independent of X ,

$$(1) \quad \mathbb{E}[(X_t - X_s)^2] = \mathbb{V}(X_1)\mathbb{E}[|t - s|] + \mathbb{E}[X_1]^2\mathbb{E}[(t - s)^2]$$

$$(2) \quad \mathbb{E}[(\bar{X}_t - \bar{X}_s)^2] \leq 16\mathbb{V}(X_1)\mathbb{E}[|t - s|] + 2(\max\{\mathbb{E}[X_1], 0\})^2\mathbb{E}[(t - s)^2],$$

[Better estimates for bounded variation case!]

- Finally, recalling that τ_{n_ℓ} is a (n_ℓ, n_ℓ) -gamma distribution with mean 1,

$$\mathbb{E}[(\tau_{n_\ell} - 1)^2] \lesssim n_\ell^{-1} \quad \text{and} \quad \mathbb{E}[|\tau_{n_\ell} - 1|] \lesssim n_\ell^{-\frac{1}{2}}.$$

- All together we get $\mathbb{V}(F^{n_\ell} - F^{n_{\ell-1}}) \lesssim n_\ell^{-\frac{1}{2}}$, and so $\beta = \frac{1}{2}$.
- Via Jensen's inequality we then easily also get $\alpha = \frac{1}{4}$.

Can we do better?

Can we do better?


- Let $n \in \mathbb{N}$ and define $\mathbf{T}(n) = \tau_{\kappa_1}$, where $\kappa_1 = \inf\{j \in \mathbb{N} : \tau_j > 1\}$. Then

$$(X_{\mathbf{T}(n)}, \bar{X}_{\mathbf{T}(n)}) \stackrel{d}{=} (V_{\kappa_1}, J_{\kappa_1})$$

where

$$V_k := \sum_{i=1}^k (S_n^i + I_n^i) \quad \text{and} \quad J_k := \bigvee_{i=1}^k (V_{i-1} + S_n^i)$$

and $\{S_n^j : j \geq 1\}$ and $\{I_n^j : j \geq 1\}$ are i.i.d. sequences of random variables with common distribution equal to \bar{X}_{e_n} and \underline{X}_{e_n} , respectively.



Can we do better?


- Let $n \in \mathbb{N}$ and define $\mathbf{T}(n) = \tau_{\kappa_1}$, where $\kappa_1 = \inf\{j \in \mathbb{N} : \tau_j > 1\}$. Then

$$(X_{\mathbf{T}(n)}, \bar{X}_{\mathbf{T}(n)}) \stackrel{d}{=} (V_{\kappa_1}, J_{\kappa_1})$$

where

$$V_k := \sum_{i=1}^k (S_n^i + I_n^i) \quad \text{and} \quad J_k := \bigvee_{i=1}^k (V_{i-1} + S_n^i)$$

and $\{S_n^j : j \geq 1\}$ and $\{I_n^j : j \geq 1\}$ are i.i.d. sequences of random variables with common distribution equal to \bar{X}_{e_n} and \underline{X}_{e_n} , respectively.



- Note that by the lack of memory property $\mathbf{T}(n) - t$ is still exponentially distributed with parameter n . Moreover

$$\mathbf{T}(n) - t \rightarrow 0 \quad \text{as } n \uparrow \infty \quad (\text{significantly faster than } \tau_n - t)$$

- $(V_{\kappa_t}, J_{\kappa_t}) \rightarrow (X_t, \bar{X}_t)$ in distribution as $n \uparrow \infty$ (rates later ...)

Can we do better?

- Now construct M independent versions of V_{κ_t} and J_{κ_t} by sampling repeatedly and indep. from the distributions of \bar{X}_{e_n} and \underline{X}_{e_n} and define

$$\widehat{F}_{MC}^{n,M} := \frac{1}{M} \sum_{i=1}^M F\left(V_{\kappa_t}^{(i)}, J_{\kappa_t}^{(i)}\right) \approx \mathbb{E}(F(X_{\mathbf{T}(n)}, \bar{X}_{\mathbf{T}(n)})) \quad (\text{sampling error})$$

$$\approx \mathbb{E}(F(X_t, \bar{X}_t)) \quad (\text{approximation error})$$

Can we do better?


- Now construct M independent versions of V_{κ_t} and J_{κ_t} by sampling repeatedly and indep. from the distributions of \bar{X}_{e_n} and \underline{X}_{e_n} and define

$$\hat{F}_{MC}^{n,M} := \frac{1}{M} \sum_{i=1}^M F\left(V_{\kappa_t}^{(i)}, J_{\kappa_t}^{(i)}\right) \approx \mathbb{E}(F(X_{\mathbf{T}(n)}, \bar{X}_{\mathbf{T}(n)})) \quad (\text{sampling error})$$

$$\approx \mathbb{E}(F(X_t, \bar{X}_t)) \quad (\text{approximation error})$$

- Levels:** Now note that $\mathbf{T}(n_{\ell-1}) \geq \mathbf{T}(n_\ell)$ and so

EITHER:



Can we do better?


- Now construct M independent versions of V_{κ_t} and J_{κ_t} by sampling repeatedly and indep. from the distributions of \bar{X}_{e_n} and \underline{X}_{e_n} and define

$$\hat{F}_{MC}^{n,M} := \frac{1}{M} \sum_{i=1}^M F(V_{\kappa_t}^{(i)}, J_{\kappa_t}^{(i)}) \approx \mathbb{E}(F(X_{\mathbf{T}(n)}, \bar{X}_{\mathbf{T}(n)})) \quad (\text{sampling error})$$

$$\approx \mathbb{E}(F(X_t, \bar{X}_t)) \quad (\text{approximation error})$$

- Levels:** Now note that $\mathbf{T}(n_{\ell-1}) \geq \mathbf{T}(n_\ell)$ and so

OR:



Can we do better?


- Now construct M independent versions of V_{κ_t} and J_{κ_t} by sampling repeatedly and indep. from the distributions of \bar{X}_{e_n} and \underline{X}_{e_n} and define

$$\hat{F}_{MC}^{n,M} := \frac{1}{M} \sum_{i=1}^M F\left(V_{\kappa_t}^{(i)}, J_{\kappa_t}^{(i)}\right) \approx \mathbb{E}(F(X_{\mathbf{T}(n)}, \bar{X}_{\mathbf{T}(n)})) \quad (\text{sampling error})$$

$$\approx \mathbb{E}(F(X_t, \bar{X}_t)) \quad (\text{approximation error})$$

- Levels:** Now note that $\mathbf{T}(n_{\ell-1}) \geq \mathbf{T}(n_\ell)$ and so

OR:



Can we do better?


- Now construct M independent versions of V_{κ_t} and J_{κ_t} by sampling repeatedly and indep. from the distributions of \bar{X}_{e_n} and \underline{X}_{e_n} and define

$$\hat{F}_{MC}^{n,M} := \frac{1}{M} \sum_{i=1}^M F(V_{\kappa_t}^{(i)}, J_{\kappa_t}^{(i)}) \approx \mathbb{E}(F(X_{\mathbf{T}(n)}, \bar{X}_{\mathbf{T}(n)})) \quad (\text{sampling error})$$

$$\approx \mathbb{E}(F(X_t, \bar{X}_t)) \quad (\text{approximation error})$$

- Levels:** Now note that $\mathbf{T}(n_{\ell-1}) \geq \mathbf{T}(n_\ell)$ and so


OR:



ETC ...

Can we do better?

EITHER, with probability $1/2$, $\mathbf{T}(n_\ell) = \mathbf{T}(n_{\ell-1})$. Hence, $F^{n_\ell} - F^{n_{\ell-1}} = 0$
 (due to the exactness of the Wiener-Hopf factorisation)
 and we have to do **no work at all** (apart from one coin toss) for this sample.




Can we do better?

OR, with probability 1/2, $\mathbf{T}(n_{\ell-1}) > \mathbf{T}(n_{\ell})$ in which case

$$\bar{X}_{\mathbf{T}(n_{\ell-1})} = \max(\bar{X}_{\mathbf{T}(n_{\ell})}, X_{\mathbf{T}(n_{\ell})} + S_{n_{\ell-1}/t})$$

$$X_{\mathbf{T}(n_{\ell-1})} = X_{\mathbf{T}(n_{\ell-1})} + S_{n_{\ell-1}/t} + I_{n_{\ell-1}/t}$$

and so the **additional work** for the path on level $\ell - 1$ is **negligible**.



Can we do better?

- Following the numerical analysis through, remembering that $\mathbf{T}(n) - 1$ is exponentially distributed, the key estimates boil down to

$$\mathbb{E}(|\mathbf{T}(n) - 1|) = \frac{1}{n} \text{ and } \mathbb{E}[(\mathbf{T}(n) - 1)^2] = \frac{1}{n^2}.$$

which imply that $\beta = 1$.

Can we do better?

- Following the numerical analysis through, remembering that $\mathbf{T}(n) - 1$ is exponentially distributed, the key estimates boil down to

$$\mathbb{E}(|\mathbf{T}(n) - 1|) = \frac{1}{n} \text{ and } \mathbb{E}[(\mathbf{T}(n) - 1)^2] = \frac{1}{n^2}.$$

which imply that $\beta = 1$.

- Back in the complexity theorem, this gives us $\alpha = \frac{1}{2}$ and $\beta = 1$ so that an expected cost of order no greater than $\mathcal{O}(\nu)$ can be delivered against an L^2 error of order $\mathcal{O}(\nu^{-\frac{1}{2}} \log^2 \nu)$. **QUASI-OPTIMAL!**

Can we do better?

- Following the numerical analysis through, remembering that $\mathbf{T}(n) - 1$ is exponentially distributed, the key estimates boil down to

$$\mathbb{E}[|\mathbf{T}(n) - 1|] = \frac{1}{n} \text{ and } \mathbb{E}[(\mathbf{T}(n) - 1)^2] = \frac{1}{n^2}.$$

which imply that $\beta = 1$.

- Back in the complexity theorem, this gives us $\alpha = \frac{1}{2}$ and $\beta = 1$ so that an expected cost of order no greater than $\mathcal{O}(\nu)$ can be delivered against an L^2 error of order $\mathcal{O}(\nu^{-\frac{1}{2}} \log^2 \nu)$. **QUASI-OPTIMAL!**
- BAD NEWS!** Because this algorithm has "awareness" of each of the exponential "Wiener-Hopf exponential periods" (i.e. we need to see whether the cumulative exponential periods exceed the level 1), it means that to implement the algorithm we need to be able to sample simultaneously from the triple $(\bar{X}_{\mathbf{e}_n}, X_{\mathbf{e}_n} - \bar{X}_{\mathbf{e}_n}, \mathbf{e}_n)$.

Comparison with other methods

Comparison with other methods

- Limited works: **[Jacod et al, 2005]** (less general – functionals of X_t only, smooth F); **[Dereich et al, 2011], [Dereich, 2011]** (more general – Lévy driven SDEs).

Comparison with other methods


- Limited works: [Jacod et al, 2005] (less general – functionals of X_t only, smooth F); [Dereich et al, 2011], [Dereich, 2011] (more general – Lévy driven SDEs).
- Rates in these papers depend on Blumenthal-Gettoor index (“jump activity”):

$$\rho := \inf \left\{ \beta > 0 : \int_{(-1,1)} |x|^\beta \Pi(dx) < \infty \right\} \in [0, 2].$$

Comparison with other methods

- Limited works: [Jacod et al, 2005] (less general – functionals of X_t only, smooth F); [Dereich et al, 2011], [Dereich, 2011] (more general – Lévy driven SDEs).
- Rates in these papers depend on Blumenthal-Gettoor index (“jump activity”):

$$\rho := \inf \left\{ \beta > 0 : \int_{(-1,1)} |x|^\beta \Pi(dx) < \infty \right\} \in [0, 2].$$



References

- Cliffe, K.A., Giles, M.B., Scheichl, R. and Teckentrup, A.L. (2011) Multilevel Monte Carlo methods and applications to elliptic PDEs with random coefficients. *Comput. Visual. Sci.* **14**(1), 3–15.
- Dereich, S. (2011) Multilevel Monte Carlo Algorithms for Lévy-driven SDEs with Gaussian correction. *Ann. Appl. Probab.* **21**(1), 283–311.
- Dereich, S. and Heidenreich, F. (2011) A multilevel Monte Carlo algorithm for Lévy-driven stochastic differential equations. *Stochastic Process. Appl.* **121**(7), 1565–1587.
- Giles, M.B. (2008) Multilevel Monte Carlo Path Simulation. *Oper. Res.* **56**(3), 607–617.
- Jacod, J., Kurtz, T.G., Méléard, S. and Protter, P. (2005) The approximate Euler method for Lévy driven stochastic differential equations. *Ann. Inst. H. Poincaré Probab. Statist.* **41**, 523–558.
- Kuznetsov (2010) Wiener-Hopf factorization and distribution of extrema for a family of Levy processes *Ann. Appl. Probab.* **20**, 1801–1830.

Kuznetsov's β -family

- The characteristic exponent ($\Psi(\theta) = -\log \mathbb{E}(e^{i\theta X_1}), \theta \in \mathbb{R}$) is given by

$$\begin{aligned} \Psi(\theta) = & \ iaz + \frac{1}{2}\sigma^2 z^2 + \frac{c_1}{\beta_1} \left\{ \mathbf{B}(\alpha_1, 1 - \lambda_1) - \mathbf{B}(\alpha_1 - \frac{i\theta}{\beta_1}, 1 - \lambda_1) \right\} \\ & + \frac{c_2}{\beta_2} \left\{ \mathbf{B}(\alpha_2, 1 - \lambda_2) - \mathbf{B}(\alpha_2 + \frac{i\theta}{\beta_2}, 1 - \lambda_2) \right\} \end{aligned}$$

where $\mathbf{B}(x, y) = \Gamma(x)\Gamma(y)/\Gamma(x + y)$ is the Beta function, with parameter range $a \in \mathbb{R}, \sigma, c_i, \alpha_i, \beta_i > 0$ and $\lambda_1, \lambda_2 \in (0, 3) \setminus \{1, 2\}$.

- The corresponding Lévy measure Π has density

$$\pi(x) = c_1 \frac{e^{-\alpha_1 \beta_1 x}}{(1 - e^{-\beta_1 x})^{\lambda_1}} \mathbf{1}_{\{x > 0\}} + c_2 \frac{e^{\alpha_2 \beta_2 x}}{(1 - e^{\beta_2 x})^{\lambda_2}} \mathbf{1}_{\{x < 0\}}.$$

The β -class of Lévy processes includes another recently introduced family of Lévy processes known as Lamperti-stable processes.

Meromorphic Lévy processes (equivalent definition)

- (i) The characteristic exponent $\Psi(z)$ is a meromorphic function which has poles at points $\{-i\rho_n, i\hat{\rho}_n\}_{n \geq 1}$, where ρ_n and $\hat{\rho}_n$ are positive real numbers.
- (ii) For $q \geq 0$ function $q + \Psi(z)$ has roots at points $\{-i\zeta_n, i\hat{\zeta}_n\}_{n \geq 1}$ where ζ_n and $\hat{\zeta}_n$ are nonnegative real numbers (strictly positive if $q > 0$). We will write $\zeta_n(q)$, $\hat{\zeta}_n(q)$ if we need to stress the dependence on q .
- (iii) The roots and poles of $q + \Psi(iz)$ satisfy the following interlacing condition

$$\dots - \rho_2 < -\zeta_2 < -\rho_1 < -\zeta_1 < 0 < \hat{\zeta}_1 < \hat{\rho}_1 < \hat{\zeta}_2 < \hat{\rho}_2 < \dots$$

- (iv) The Wiener-Hopf factors are expressed as convergent infinite products,

$$\mathbb{E} \left[e^{-z\bar{X}_{e_q}} \right] = \prod_{n \geq 1} \frac{1 + \frac{z}{\rho_n}}{1 + \frac{z}{\zeta_n}}$$

$$\mathbb{E} \left[e^{zX_{e_q}} \right] = \prod_{n \geq 1} \frac{1 + \frac{z}{\hat{\rho}_n}}{1 + \frac{z}{\hat{\zeta}_n}}.$$

Distribution of extrema

- For $x \geq 0$

$$\mathbb{P}(\overline{X}_{e_q} \in dx) = a_0(\rho, \zeta)\delta_0(dx) + \sum_{n=1}^{\infty} a_n(\rho, \zeta)\zeta_n e^{-\zeta_n x} dx$$

- Here

$$a_0(\rho, \zeta) = \lim_{n \rightarrow +\infty} \prod_{k=1}^n \frac{\zeta_k}{\rho_k}, \quad a_n(\rho, \zeta) = \left(1 - \frac{\zeta_n}{\rho_n}\right) \prod_{\substack{k \geq 1 \\ k \neq n}} \frac{1 - \frac{\zeta_n}{\rho_k}}{1 - \frac{\zeta_n}{\zeta_k}}$$

- A similar expression holds for $\mathbb{P}(-\underline{X}_{e_q} \in dx)$.