

# A Ciesielski-Taylor type identity for positive self-similar Markov processes

**A. E. Kyprianou** and **P. Patie**

Department of Mathematical Sciences, University of Bath

## Classical Ciesielski-Taylor Identity

## Classical Ciesielski-Taylor Identity

- $(X, Q^{(\nu)})$  is a Bessel process  $X$  starting from 0 with dimension  $\nu > 0$ .

## Classical Ciesielski-Taylor Identity

- $(X, Q^{(\nu)})$  is a Bessel process  $X$  starting from 0 with dimension  $\nu > 0$ .
- i.e. the  $[0, \infty)$ -valued diffusion whose infinitesimal generator is given by

$$L_\nu f(x) = \frac{1}{2}f''(x) + \frac{\nu - 1}{2x}f'(x)$$

on  $(0, \infty)$  for  $f \in C^2(0, \infty)$  with instantaneous reflection at 0 when  $\nu \in (0, 2)$  (i.e.  $f'(0^+) = 0$ ) and when  $\nu \geq 2$  the origin is an entrance-non-exit boundary point.

## Classical Ciesielski-Taylor Identity

- $(X, Q^{(\nu)})$  is a Bessel process  $X$  starting from 0 with dimension  $\nu > 0$ .
- i.e. the  $[0, \infty)$ -valued diffusion whose infinitesimal generator is given by

$$L_\nu f(x) = \frac{1}{2}f''(x) + \frac{\nu - 1}{2x}f'(x)$$

on  $(0, \infty)$  for  $f \in C^2(0, \infty)$  with instantaneous reflection at 0 when  $\nu \in (0, 2)$  (i.e.  $f'(0^+) = 0$ ) and when  $\nu \geq 2$  the origin is an entrance-non-exit boundary point.

- Ciesielski and Taylor (1962) and later Gettoor and Sharp (1979) show: For  $a > 0$  and any  $\nu > 0$ ,

$$\left(T_a, Q^{(\nu)}\right) \stackrel{(d)}{=} \left(\int_0^\infty \mathbb{I}_{\{X_s \leq a\}} ds, Q^{(\nu+2)}\right) \quad (1)$$

where

$$T_a = \inf\{s \geq 0; X_s = a\}.$$

## Subsequent work on Ciesielski-Taylor Identities

## Subsequent work on Ciesielski-Taylor Identities

- Biane (1985) Gives a generalisation to one dimensional diffusions.

## Subsequent work on Ciesielski-Taylor Identities

- Biane (1985) Gives a generalisation to one dimensional diffusions.
- Yor (1991) gives a probabilistic explanation of the original C-T identity through occupation time formula and Ray-Knight theorems.



## Subsequent work on Ciesielski-Taylor Identities

- Biane (1985) Gives a generalisation to one dimensional diffusions.
- Yor (1991) gives a probabilistic explanation of the original C-T identity through occupation time formula and Ray-Knight theorems.
- Carmona, Petit and Yor (1998) establish a similar identity for positive self-similar 'saw-tooth' process.

## Subsequent work on Ciesielski-Taylor Identities

- Biane (1985) Gives a generalisation to one dimensional diffusions.
- Yor (1991) gives a probabilistic explanation of the original C-T identity through occupation time formula and Ray-Knight theorems.
- Carmona, Petit and Yor (1998) establish a similar identity for positive self-similar 'saw-tooth' process.
- Bertoin (1992) remarks that a C-T-type identity can be observed in the setting of a reflected spectrally negative stable Lévy process.

## Subsequent work on Ciesielski-Taylor Identities

- Biane (1985) Gives a generalisation to one dimensional diffusions.
- Yor (1991) gives a probabilistic explanation of the original C-T identity through occupation time formula and Ray-Knight theorems.
- Carmona, Petit and Yor (1998) establish a similar identity for positive self-similar 'saw-tooth' process.
- Bertoin (1992) remarks that a C-T-type identity can be observed in the setting of a reflected spectrally negative stable Lévy process.

One begins to get the whiff of the possibility of a general result for positive self-similar Markov processes.

## Positive self-similar Markov process

## Positive self-similar Markov process

- $\alpha$ -pssMp: A  $[0, \infty)$ -valued Feller process which enjoys the following  $\alpha$ -self-similarity property, where  $\alpha > 0$ . For any  $x > 0$ , and  $c > 0$ ,

$$((X_t)_{t \geq 0}, \mathbb{P}_{cx}) \stackrel{(d)}{=} ((cX_{c^{-\alpha}t})_{t \geq 0}, \mathbb{P}_x).$$

## Positive self-similar Markov process

- $\alpha$ -pssMp: A  $[0, \infty)$ -valued Feller process which enjoys the following  $\alpha$ -self-similarity property, where  $\alpha > 0$ . For any  $x > 0$ , and  $c > 0$ ,

$$((X_t)_{t \geq 0}, P_{cx}) \stackrel{(d)}{=} ((cX_{c^{-\alpha}t})_{t \geq 0}, P_x).$$

- Lamperti (1972) showed that, for any  $x \in \mathbb{R}$ , there exists a one to one mapping between  $\mathbb{P}_x$ , the law of a generic Lévy process (possibly killed at an independent and exponentially distributed time), say  $\xi = (\xi_t : t \geq 0)$ , starting from  $x$ , and the law  $P_{e^x}$  via the relation

$$X_t = e^{\xi_{A_t}}, \quad 0 \leq t < \zeta,$$

where  $\zeta = \inf\{t > 0 : X_t = 0\}$  and

$$A_t = \inf\{s \geq 0; \int_0^s e^{\alpha \xi_u} du > t\}.$$

## Spectrally negative, entrance law, recurrent extension

## Spectrally negative, entrance law, recurrent extension

- In this talk we are interested in the case that  $X$  (and hence  $\xi$ ) has only negative jumps.



## Spectrally negative, entrance law, recurrent extension

- In this talk we are interested in the case that  $X$  (and hence  $\xi$ ) has only negative jumps.
- When  $\mathbb{E}(\xi_1) \geq 0$  and  $\xi$  is not killed, one may extend the definition of  $X$  to include the case that it is issued from the origin by establishing its entrance law  $P_0$  as the weak limit with respect to the Skorohod topology of  $P_x$  as  $x \downarrow 0$ . Bertoin and Yor (2002).

## Spectrally negative, entrance law, recurrent extension

- In this talk we are interested in the case that  $X$  (and hence  $\xi$ ) has only negative jumps.
- When  $\mathbb{E}(\xi_1) \geq 0$  and  $\xi$  is not killed, one may extend the definition of  $X$  to include the case that it is issued from the origin by establishing its entrance law  $P_0$  as the weak limit with respect to the Skorohod topology of  $P_x$  as  $x \downarrow 0$ . Bertoin and Yor (2002).
- When  $\mathbb{E}(\xi_1) < 0$  (resp.  $\xi$  is killed) then the boundary state 0 is reached continuously (resp. by a jump). In these two cases, one cannot construct an entrance law, however, Rivero (2005) and Fitzsimmons (2006), show that it is possible instead to construct a unique recurrent extension on  $[0, \infty)$  such that paths leave 0 continuously, thereby giving a meaning to  $P_0$ , if and only if there exists a  $\theta \in (0, \alpha)$  such that  $\mathbb{E}(e^{\theta \xi_1}) = 1$ .

## Spectrally negative, entrance law, recurrent extension

- In this talk we are interested in the case that  $X$  (and hence  $\xi$ ) has only negative jumps.
- When  $\mathbb{E}(\xi_1) \geq 0$  and  $\xi$  is not killed, one may extend the definition of  $X$  to include the case that it is issued from the origin by establishing its entrance law  $P_0$  as the weak limit with respect to the Skorohod topology of  $P_x$  as  $x \downarrow 0$ . Bertoin and Yor (2002).
- When  $\mathbb{E}(\xi_1) < 0$  (resp.  $\xi$  is killed) then the boundary state 0 is reached continuously (resp. by a jump). In these two cases, one cannot construct an entrance law, however, Rivero (2005) and Fitzsimmons (2006), show that it is possible instead to construct a unique recurrent extension on  $[0, \infty)$  such that paths leave 0 continuously, thereby giving a meaning to  $P_0$ , if and only if there exists a  $\theta \in (0, \alpha)$  such that  $\mathbb{E}(e^{\theta \xi_1}) = 1$ .
- **Objective of this talk: Fix  $\alpha > 0$  and show that for a given spectrally negative Lévy process fitting the previous two categories, and hence given the associated law  $P_0$ ,**

$$(T_a, P_0) \stackrel{(d)}{=} \left( \int_0^\infty \mathbb{I}_{\{X_s \leq a\}} ds, P_0^* \right),$$

where  $T_a = \inf\{t > 0 : X_t = a\}$  and  $P_0^*$  is to be identified.

## A new transformation for spectrally negative Lévy processes: $\mathcal{T}_\beta$

## A new transformation for spectrally negative Lévy processes: $\mathcal{T}_\beta$

- For any spectrally negative Lévy process, henceforth denoted by  $\xi = (\xi_t, t \geq 0)$ , we define simultaneously the Laplace exponent and its law  $\mathbb{P}^\psi$  by

$$\psi(u) = \log \mathbb{E}^\psi(\exp\{u\xi_1\}), \quad u \geq 0.$$

**Warning: included in the definition of  $\xi$  is the possibility of exponential killing!**

## A new transformation for spectrally negative Lévy processes: $\mathcal{T}_\beta$

- For any spectrally negative Lévy process, henceforth denoted by  $\xi = (\xi_t, t \geq 0)$ , we define simultaneously the Laplace exponent and its law  $\mathbb{P}^\psi$  by

$$\psi(u) = \log \mathbb{E}^\psi(\exp\{u\xi_1\}), \quad u \geq 0.$$

**Warning: included in the definition of  $\xi$  is the possibility of exponential killing!**

- For each fixed  $\beta > 0$ , define the linear transformation

$$\mathcal{T}_\beta\psi(u) = \frac{u}{u + \beta}\psi(u + \beta), \quad u \geq 0.$$

## A new transformation for spectrally negative Lévy processes: $\mathcal{T}_\beta$

- For any spectrally negative Lévy process, henceforth denoted by  $\xi = (\xi_t, t \geq 0)$ , we define simultaneously the Laplace exponent and its law  $\mathbb{P}^\psi$  by

$$\psi(u) = \log \mathbb{E}^\psi(\exp\{u\xi_1\}), \quad u \geq 0.$$

**Warning: included in the definition of  $\xi$  is the possibility of exponential killing!**

- For each fixed  $\beta > 0$ , define the linear transformation

$$\mathcal{T}_\beta\psi(u) = \frac{u}{u + \beta}\psi(u + \beta), \quad u \geq 0.$$

- **Theorem:**  $\mathcal{T}_\beta\psi$  is the Laplace exponent of another spectrally negative Lévy process with no exponential killing.

## A new transformation for spectrally negative Lévy processes: $\mathcal{T}_\beta$

- For any spectrally negative Lévy process, henceforth denoted by  $\xi = (\xi_t, t \geq 0)$ , we define simultaneously the Laplace exponent and its law  $\mathbb{P}^\psi$  by

$$\psi(u) = \log \mathbb{E}^\psi(\exp\{u\xi_1\}), \quad u \geq 0.$$

**Warning: included in the definition of  $\xi$  is the possibility of exponential killing!**

- For each fixed  $\beta > 0$ , define the linear transformation

$$\mathcal{T}_\beta\psi(u) = \frac{u}{u + \beta}\psi(u + \beta), \quad u \geq 0.$$

- Theorem:**  $\mathcal{T}_\beta\psi$  is the Laplace exponent of another spectrally negative Lévy process with no exponential killing.
- Easy proof when  $\psi$  has no killing and  $\psi'(0+) \geq 0$ :**

Algebra:  $\mathcal{T}_\beta\psi(u) = \mathcal{E}_\beta\psi(u) - \beta\mathcal{E}_\beta\phi(u)$

$\mathcal{E}_\beta f(u) = f(u + \beta) - f(\beta)$  (Esscher transform)

$\phi(u) = \psi(u)/u$  (Laplace exponent of descending ladder height subordinator).



## Ciesielski-Taylor identity for spectrally negative pssMp

### Theorem

Fix  $\alpha > 0$ . Suppose that  $\psi$  is the Laplace exponent of a spectrally negative Lévy process. Assume that  $\theta$ , the largest root in  $[0, \infty)$  of the equation  $\psi(\theta) = 0$ , satisfies  $\theta < \alpha$ . Let  $\mathbb{P}_0^\psi$  be the law of the pssMp associated with  $\mathbb{P}^\psi$  and issued from 0. Then for any  $a > 0$ , the following Ciesielski-Taylor type identity in law

$$\left( T_a, \mathbb{P}_0^\psi \right) \stackrel{(d)}{=} \left( \int_0^\infty \mathbb{I}_{\{X_s \leq a\}} ds, \mathbb{P}_0^{\mathcal{T}_{\alpha\psi}} \right)$$

holds.

## A fluctuation identity of Patie

## A fluctuation identity of Patie

- Assume that  $0 \leq \theta < \alpha$  where  $\psi(\theta) = 0$ .

## A fluctuation identity of Patie

- Assume that  $0 \leq \theta < \alpha$  where  $\psi(\theta) = 0$ .
- Let

$$a_n(\psi; \alpha)^{-1} = \prod_{k=1}^n \psi(\alpha k), \quad a_0 = 1,$$

## A fluctuation identity of Patie

- Assume that  $0 \leq \theta < \alpha$  where  $\psi(\theta) = 0$ .
- Let

$$a_n(\psi; \alpha)^{-1} = \prod_{k=1}^n \psi(\alpha k), \quad a_0 = 1,$$

- Define the positive and entire function  $\mathcal{I}_{\psi, \alpha}(z)$  which admits the series representation

$$\mathcal{I}_{\psi, \alpha}(z) = \sum_{n=0}^{\infty} a_n(\psi; \alpha) z^n, \quad z \in \mathbb{C}.$$

## A fluctuation identity of Patie

- Assume that  $0 \leq \theta < \alpha$  where  $\psi(\theta) = 0$ .
- Let

$$a_n(\psi; \alpha)^{-1} = \prod_{k=1}^n \psi(\alpha k), \quad a_0 = 1,$$

- Define the positive and entire function  $\mathcal{I}_{\psi, \alpha}(z)$  which admits the series representation

$$\mathcal{I}_{\psi, \alpha}(z) = \sum_{n=0}^{\infty} a_n(\psi; \alpha) z^n, \quad z \in \mathbb{C}.$$

- Theorem (Patie):** For  $0 \leq x \leq a$  and  $q \geq 0$ , we have

$$\mathbb{E}_x^\psi \left[ e^{-qT_a} \right] = \frac{\mathcal{I}_{\psi, \alpha}(qx^\alpha)}{\mathcal{I}_{\psi, \alpha}(qa^\alpha)}.$$

In particular

$$\mathbb{E}_0^\psi \left[ e^{-qT_a} \right] = \frac{1}{\mathcal{I}_{\psi, \alpha}(qa^\alpha)}.$$

## How the proof works: I

## How the proof works: I

- Define

$$O_q^{\mathcal{T}\alpha\psi}(x; a) = \mathbb{E}_x^{\mathcal{T}\alpha\psi} \left[ e^{-q \int_0^\infty \mathbb{I}_{\{X_s \leq a\}} ds} \right]$$

and try to show that  $O_q^{\mathcal{T}\alpha\psi}(0; a) = \mathbb{E}_0^\psi [e^{-qT_a}] = 1/\mathcal{I}_{\psi, \alpha}(qa^\alpha)$  for all  $q \geq 0$ .



## How the proof works: I

- Define

$$O_q^{\mathcal{T}_{\alpha}\psi}(x; a) = \mathbb{E}_x^{\mathcal{T}_{\alpha}\psi} \left[ e^{-q \int_0^{\infty} \mathbb{I}_{\{X_s \leq a\}} ds} \right]$$

and try to show that  $O_q^{\mathcal{T}_{\alpha}\psi}(0; a) = \mathbb{E}_0^{\psi} [e^{-qT_a}] = 1/\mathcal{I}_{\psi, \alpha}(qa^{\alpha})$  for all  $q \geq 0$ .

- Self-similarity of  $X$  means that

$$O_q^{\mathcal{T}_{\alpha}\psi}(0; a) = O_{qa^{\alpha}}^{\mathcal{T}_{\alpha}\psi}(0; 1),$$

so enough to establish the suggested equality when  $a = 1$ .

## How the proof works: I

- Define

$$O_q^{\mathcal{T}_{\alpha}\psi}(x; a) = \mathbb{E}_x^{\mathcal{T}_{\alpha}\psi} \left[ e^{-q \int_0^{\infty} \mathbb{I}_{\{X_s \leq a\}} ds} \right]$$

and try to show that  $O_q^{\mathcal{T}_{\alpha}\psi}(0; a) = \mathbb{E}_0^{\psi} [e^{-qT_a}] = 1/\mathcal{I}_{\psi, \alpha}(qa^{\alpha})$  for all  $q \geq 0$ .

- Self-similarity of  $X$  means that

$$O_q^{\mathcal{T}_{\alpha}\psi}(0; a) = O_{qa^{\alpha}}^{\mathcal{T}_{\alpha}\psi}(0; 1),$$

so enough to establish the suggested equality when  $a = 1$ .

- Next, note using spectral negativity,

$$O_q^{\mathcal{T}_{\alpha}\psi}(0; 1) = \mathbb{E}_0^{\mathcal{T}_{\alpha}\psi}(e^{-qT_1}) O_q^{\mathcal{T}_{\alpha}\psi}(1; 1)$$

so it is enough to show that

$$O_q^{\mathcal{T}_{\alpha}\psi}(1; 1) = \frac{\mathcal{I}_{\mathcal{T}_{\alpha}\psi, \alpha}(q)}{\mathcal{I}_{\psi, \alpha}(q)}.$$

## How the proof works: II

## How the proof works: II

- Let  $\tau_1 = \inf\{s > 0; X_s < 1\}$  and  $T_1 = \inf\{s > 0 : X_s = 1\}$ .

## How the proof works: II

- Let  $\tau_1 = \inf\{s > 0; X_s < 1\}$  and  $T_1 = \inf\{s > 0 : X_s = 1\}$ .
- Fix  $y > 1$ .

## How the proof works: II

- Let  $\tau_1 = \inf\{s > 0; X_s < 1\}$  and  $T_1 = \inf\{s > 0 : X_s = 1\}$ .
- Fix  $y > 1$ .
- Use Strong Markov Property,  $(\mathcal{T}_\alpha \psi)'(0^+) = \psi(\alpha)/\alpha > 0$  and spectral negativity:

$$\begin{aligned}
 & O_q^{\mathcal{T}_\alpha \psi}(1; 1) \\
 = & \mathbb{E}_1^{\mathcal{T}_\alpha \psi} \left[ e^{-q \int_0^{T_y} \mathbb{1}_{\{X_s \leq 1\}} ds} \right] \left( \mathbb{E}_y^{\mathcal{T}_\alpha \psi} [\mathbb{1}_{\{\tau_1 < \infty\}}] \mathbb{E}_{X_{\tau_1}}^{\mathcal{T}_\alpha \psi} [e^{-qT_1}] \right) O_q^{\mathcal{T}_\alpha \psi}(1; 1) + \mathbb{E}_y^{\mathcal{T}_\alpha \psi} [\mathbb{1}_{\{\tau_1 = \infty\}}]
 \end{aligned}$$

## How the proof works: II

- Let  $\tau_1 = \inf\{s > 0; X_s < 1\}$  and  $T_1 = \inf\{s > 0 : X_s = 1\}$ .
- Fix  $y > 1$ .
- Use Strong Markov Property,  $(\mathcal{T}_\alpha \psi)'(0^+) = \psi(\alpha)/\alpha > 0$  and spectral negativity:

$$O_q^{\mathcal{T}_\alpha \psi}(1; 1) = \mathbf{E}_1^{\mathcal{T}_\alpha \psi} \left[ e^{-q \int_0^{T_y} \mathbb{1}_{\{X_s \leq 1\}} ds} \right] \left( \mathbf{E}_y^{\mathcal{T}_\alpha \psi} [\mathbb{I}_{\{\tau_1 < \infty\}}] \mathbf{E}_{X_{\tau_1}}^{\mathcal{T}_\alpha \psi} [e^{-q T_1}] \right) O_q^{\mathcal{T}_\alpha \psi}(1; 1) + \mathbf{E}_y^{\mathcal{T}_\alpha \psi} [\mathbb{I}_{\{\tau_1 = \infty\}}]$$

- Solving for  $O_q^{\mathcal{T}_\alpha \psi}(1; 1)$  we get

$$O_q^{\mathcal{T}_\alpha \psi}(1; 1) = \frac{\mathbf{E}_y^{\mathcal{T}_\alpha \psi} [\mathbb{I}_{\{\tau_1 = \infty\}}]}{\left\{ \mathbf{E}_1^{\mathcal{T}_\alpha \psi} \left[ e^{-q \int_0^{T_y} \mathbb{1}_{\{X_s \leq 1\}} ds} \right] \right\}^{-1} - \mathbf{E}_y^{\mathcal{T}_\alpha \psi} [\mathbb{I}_{\{\tau_1 < \infty\}}] \mathbf{E}_{X_{\tau_1}}^{\mathcal{T}_\alpha \psi} [e^{-q T_1}]}$$

## How the proof works: II

- Let  $\tau_1 = \inf\{s > 0; X_s < 1\}$  and  $T_1 = \inf\{s > 0 : X_s = 1\}$ .
- Fix  $y > 1$ .
- Use Strong Markov Property,  $(\mathcal{T}_\alpha\psi)'(0^+) = \psi(\alpha)/\alpha > 0$  and spectral negativity:

$$O_q^{\mathcal{T}_\alpha\psi}(1; 1) = \mathbf{E}_1^{\mathcal{T}_\alpha\psi} \left[ e^{-q \int_0^{T_y} \mathbb{1}_{\{X_s \leq 1\}} ds} \right] \left( \mathbf{E}_y^{\mathcal{T}_\alpha\psi} [\mathbb{I}_{\{\tau_1 < \infty\}}] \mathbf{E}_{X_{\tau_1}}^{\mathcal{T}_\alpha\psi} [e^{-qT_1}] \right) O_q^{\mathcal{T}_\alpha\psi}(1; 1) + \mathbf{E}_y^{\mathcal{T}_\alpha\psi} [\mathbb{I}_{\{\tau_1 = \infty\}}]$$

- Solving for  $O_q^{\mathcal{T}_\alpha\psi}(1; 1)$  we get

$$O_q^{\mathcal{T}_\alpha\psi}(1; 1) = \frac{\mathbf{E}_y^{\mathcal{T}_\alpha\psi} [\mathbb{I}_{\{\tau_1 = \infty\}}]}{\left\{ \mathbf{E}_1^{\mathcal{T}_\alpha\psi} \left[ e^{-q \int_0^{T_y} \mathbb{1}_{\{X_s \leq 1\}} ds} \right] \right\}^{-1} - \mathbf{E}_y^{\mathcal{T}_\alpha\psi} [\mathbb{I}_{\{\tau_1 < \infty\}}] \mathbf{E}_{X_{\tau_1}}^{\mathcal{T}_\alpha\psi} [e^{-qT_1}]}$$

- Proof is formalised by taking limits as  $y \downarrow 1$ .



## How the proof works III: special case, bounded variation paths

## How the proof works III: special case, bounded variation paths

- Let  $\tau_0^\xi = \inf\{t > 0 : \xi_t < 0\}$ . Bounded variation paths for  $\xi$  implies that  $\mathbb{P}^{\mathcal{T}_\alpha^\psi}(\tau_0^\xi > 0) = 1$ .

## How the proof works III: special case, bounded variation paths

- Let  $\tau_0^\xi = \inf\{t > 0 : \xi_t < 0\}$ . Bounded variation paths for  $\xi$  implies that  $\mathbb{P}^{\mathcal{T}_\alpha^\psi}(\tau_0^\xi > 0) = 1$ .
- Hence  $\mathbb{P}_1^{\mathcal{T}_\alpha^\psi}(\tau_1 > 0) = 1$ . Hence can just set  $y = 0$  in the formula for  $O_q^{\mathcal{T}_\alpha^\psi}(1; 1)$ .

## How the proof works III: special case, bounded variation paths

- Let  $\tau_0^\xi = \inf\{t > 0 : \xi_t < 0\}$ . Bounded variation paths for  $\xi$  implies that  $\mathbb{P}^{\mathcal{T}_\alpha^\psi}(\tau_0^\xi > 0) = 1$ .
- Hence  $\mathbb{P}_1^{\mathcal{T}_\alpha^\psi}(\tau_1 > 0) = 1$ . Hence can just set  $y = 0$  in the formula for  $O_q^{\mathcal{T}_\alpha^\psi}(1; 1)$ .
- In which case,  $\mathbb{E}_1^{\mathcal{T}_\alpha^\psi} \left[ e^{-q \int_0^{T_1} \mathbb{1}_{\{X_s \leq 1\}} ds} \right] = 1$ .

## How the proof works III: special case, bounded variation paths

- Let  $\tau_0^\xi = \inf\{t > 0 : \xi_t < 0\}$ . Bounded variation paths for  $\xi$  implies that  $\mathbb{P}^{\mathcal{T}_\alpha\psi}(\tau_0^\xi > 0) = 1$ .
- Hence  $\mathbb{P}_1^{\mathcal{T}_\alpha\psi}(\tau_1 > 0) = 1$ . Hence can just set  $y = 0$  in the formula for  $O_q^{\mathcal{T}_\alpha\psi}(1; 1)$ .
- In which case,  $\mathbb{E}_1^{\mathcal{T}_\alpha\psi} \left[ e^{-q \int_0^{T_1} \mathbb{1}_{\{X_s \leq 1\}} ds} \right] = 1$ .
- On the one hand, recalling that  $(\mathcal{T}_\alpha\psi)'(0^+) = \psi(\alpha)/\alpha > 0$ , we observe that

$$\mathbb{E}_1^{\mathcal{T}_\alpha\psi} [\mathbb{1}_{\{\tau_1 = \infty\}}] = \mathbb{P}_0^{\mathcal{T}_\alpha\psi} (\tau_0^\xi = \infty) = \frac{1}{(\mathcal{T}_\alpha\psi)'(0^+)} W_{\mathcal{T}_\alpha\psi}(0^+) = \frac{\psi(\alpha)}{\alpha} W_{\mathcal{T}_\alpha\psi}(0^+) > 0$$

Here  $W_{\mathcal{T}_\alpha\psi}$  is the scale function under  $\mathbb{P}^{\mathcal{T}_\alpha\psi}$ . In other words it is the unique continuous function on  $[0, \infty)$  whose Laplace transform satisfies

$$\int_0^\infty e^{-ux} W_{\mathcal{T}_\alpha\psi}(x) dx = \frac{1}{\mathcal{T}_\alpha\psi(u)}.$$

## How the proof works IV: special case, bounded variation paths

## How the proof works IV: special case, bounded variation paths

- For the remaining term, by Fubini's theorem (recalling the positivity of coefficients in the definition of  $\mathcal{I}_{\mathcal{T}_\alpha\psi, \alpha}$ ), we have with the help of Patie's identity,

$$\begin{aligned}
 & \mathbb{E}_1^{\mathcal{T}_\alpha\psi} \left[ \mathbb{I}_{\{\tau_1 < \infty\}} \mathbb{E}_{X_{\tau_1}^{\mathcal{T}_\alpha\psi}} \left[ e^{-qT_1} \right] \right] \\
 &= \mathbb{E}_1^{\mathcal{T}_\alpha\psi} \left[ \frac{\mathcal{I}_{\mathcal{T}_\alpha\psi, \alpha}(qX_{\tau_1}^\alpha) \mathbb{I}_{\{\tau_1 < \infty\}}}{\mathcal{I}_{\mathcal{T}_\alpha\psi, \alpha}(q)} \right] \\
 &= \frac{1}{\mathcal{I}_{\mathcal{T}_\alpha\psi, \alpha}(q)} \mathbb{E}_1^{\mathcal{T}_\alpha\psi} \left[ \sum_{n=0}^{\infty} a_n(\mathcal{T}_\alpha\psi; \alpha) q^n X_{\tau_1}^{\alpha n} \mathbb{I}_{\{\tau_1 < \infty\}} \right] \\
 &= \frac{1}{\mathcal{I}_{\mathcal{T}_\alpha\psi, \alpha}(q)} \sum_{n=0}^{\infty} a_n(\mathcal{T}_\alpha\psi; \alpha) q^n \mathbb{E}_0^{\mathcal{T}_\alpha\psi} \left[ e^{\alpha n \xi_{\tau_0}^\xi} \mathbb{I}_{\{\tau_0^\xi < \infty\}} \right].
 \end{aligned}$$

## How the proof works IV: special case, bounded variation paths

- For the remaining term, by Fubini's theorem (recalling the positivity of coefficients in the definition of  $\mathcal{I}_{\mathcal{T}_\alpha\psi, \alpha}$ ), we have with the help of Patie's identity,

$$\begin{aligned}
 & \mathbb{E}_1^{\mathcal{T}_\alpha\psi} \left[ \mathbb{I}_{\{\tau_1 < \infty\}} \mathbb{E}_{X_{\tau_1}}^{\mathcal{T}_\alpha\psi} \left[ e^{-qT_1} \right] \right] \\
 &= \mathbb{E}_1^{\mathcal{T}_\alpha\psi} \left[ \frac{\mathcal{I}_{\mathcal{T}_\alpha\psi, \alpha}(qX_{\tau_1}^\alpha) \mathbb{I}_{\{\tau_1 < \infty\}}}{\mathcal{I}_{\mathcal{T}_\alpha\psi, \alpha}(q)} \right] \\
 &= \frac{1}{\mathcal{I}_{\mathcal{T}_\alpha\psi, \alpha}(q)} \mathbb{E}_1^{\mathcal{T}_\alpha\psi} \left[ \sum_{n=0}^{\infty} a_n(\mathcal{T}_\alpha\psi; \alpha) q^n X_{\tau_1}^{\alpha n} \mathbb{I}_{\{\tau_1 < \infty\}} \right] \\
 &= \frac{1}{\mathcal{I}_{\mathcal{T}_\alpha\psi, \alpha}(q)} \sum_{n=0}^{\infty} a_n(\mathcal{T}_\alpha\psi; \alpha) q^n \mathbb{E}_0^{\mathcal{T}_\alpha\psi} \left[ e^{\alpha n \xi_{\tau_0}^\xi} \mathbb{I}_{\{\tau_0^\xi < \infty\}} \right].
 \end{aligned}$$

- Classical fluctuation theory for spectrally negative Lévy processes gives (in the bounded variation case):

$$\mathbb{E}_0^{\mathcal{T}_\alpha\psi} \left( e^{u \xi_{\tau_0}^\xi} \mathbb{I}_{\{\tau_0^\xi < \infty\}} \right) = 1 - \frac{\mathcal{T}_\alpha\psi(u)}{u} W_{\mathcal{T}_\alpha\psi}(0+).$$



## How the proof works V:

## How the proof works V:

- Putting the bits together

$$\begin{aligned}
 & O_q^{\mathcal{T}_\alpha \psi}(1; 1) \\
 &= \frac{\psi(\alpha) W_{\mathcal{T}_\alpha \psi}(0+)/\alpha}{1 - \frac{\sum_{n=0}^{\infty} a_n(\mathcal{T}_\alpha \psi; \alpha) q^n \left\{ 1 - \frac{\mathcal{T}_\alpha \psi(\alpha n)}{\alpha n} y^{-\alpha n} W_{\mathcal{T}_\alpha \psi}(0+) \right\}}{\mathcal{I}_{\mathcal{T}_\alpha \psi, \alpha}(q)}} \\
 &= \frac{\psi(\alpha) \mathcal{I}_{\mathcal{T}_\alpha \psi, \alpha}(q)}{\alpha \sum_{n=0}^{\infty} a_n(\mathcal{T}_\alpha \psi; \alpha) q^n \frac{\mathcal{T}_\alpha \psi(\alpha n)}{\alpha n}}.
 \end{aligned}$$

## How the proof works V:

- Putting the bits together

$$\begin{aligned}
 O_q^{\mathcal{T}_\alpha \psi}(1; 1) &= \frac{\psi(\alpha) W_{\mathcal{T}_\alpha \psi}(0+)/\alpha}{1 - \frac{\sum_{n=0}^{\infty} a_n(\mathcal{T}_\alpha \psi; \alpha) q^n \left\{ 1 - \frac{\mathcal{T}_\alpha \psi(\alpha n)}{\alpha n} y^{-\alpha n} W_{\mathcal{T}_\alpha \psi}(0+) \right\}}{\mathcal{I}_{\mathcal{T}_\alpha \psi, \alpha}(q)}} \\
 &= \frac{\psi(\alpha) \mathcal{I}_{\mathcal{T}_\alpha \psi, \alpha}(q)}{\alpha \sum_{n=0}^{\infty} a_n(\mathcal{T}_\alpha \psi; \alpha) q^n \frac{\mathcal{T}_\alpha \psi(\alpha n)}{\alpha n}}.
 \end{aligned}$$

- Next, observing that, for any  $n \geq 1$ ,

$$\frac{\psi(\alpha) \alpha n}{\alpha \mathcal{T}_\alpha \psi(\alpha n)} a_n(\mathcal{T}_\alpha \psi; \alpha)^{-1} = \prod_{k=1}^n \psi(\alpha k) \quad (= 1 \text{ when } n = 0).$$

we deduce, as required, the identity

$$O_q^{\mathcal{T}_\alpha \psi}(1; 1) = \frac{\mathcal{I}_{\mathcal{T}_\alpha \psi, \alpha}(q)}{\sum_{n=0}^{\infty} a_n(\psi; \alpha) q^n} = \frac{\mathcal{I}_{\mathcal{T}_\alpha \psi, \alpha}(q)}{\mathcal{I}_{\psi, \alpha}(q)}.$$

## Example: cross reference against the original CT identity

## Example: cross reference against the original CT identity

- When  $\psi$  has no jump component it is possible to extract the original Ciesielski-Taylor identity for Bessel processes.

## Example: cross reference against the original CT identity

- When  $\psi$  has no jump component it is possible to extract the original Ciesielski-Taylor identity for Bessel processes.
- Indeed, we may take  $\alpha = 2$  and

$$\psi_\nu(u) = \frac{1}{2}u^2 + \left(\frac{\nu}{2} - 1\right)u$$

where  $\nu > 0$ . In that case it follows that  $P^{\psi_\nu}$  is the law of a Bessel process of dimension  $\nu$  as described in the introduction.

## Example: cross reference against the original CT identity

- When  $\psi$  has no jump component it is possible to extract the original Ciesielski-Taylor identity for Bessel processes.
- Indeed, we may take  $\alpha = 2$  and

$$\psi_\nu(u) = \frac{1}{2}u^2 + \left(\frac{\nu}{2} - 1\right)u$$

where  $\nu > 0$ . In that case it follows that  $P^{\psi_\nu}$  is the law of a Bessel process of dimension  $\nu$  as described in the introduction.

- Note that the root  $\theta$  is zero for  $\nu \geq 2$  and when  $\nu \in (0, 2)$  we have  $\theta = 2 - \nu < 2 = \alpha$  thereby fulfilling the required condition of the main result.

## Example: cross reference against the original CT identity

- When  $\psi$  has no jump component it is possible to extract the original Ciesielski-Taylor identity for Bessel processes.
- Indeed, we may take  $\alpha = 2$  and

$$\psi_\nu(u) = \frac{1}{2}u^2 + \left(\frac{\nu}{2} - 1\right)u$$

where  $\nu > 0$ . In that case it follows that  $P^{\psi_\nu}$  is the law of a Bessel process of dimension  $\nu$  as described in the introduction.

- Note that the root  $\theta$  is zero for  $\nu \geq 2$  and when  $\nu \in (0, 2)$  we have  $\theta = 2 - \nu < 2 = \alpha$  thereby fulfilling the required condition of the main result.
- The transformation  $\mathcal{T}_2$  gives us the new Laplace exponent

$$\mathcal{T}_2\psi_\nu(u) = \frac{1}{2}u^2 + \frac{\nu}{2}u = \psi_{\nu+2}(u).$$